

HEMEXOL

VIRUCIDAL POWDER

HEMEXOL VIRUCIDAL POWDER

Based on a stabilized blend of peroxygen compounds. Effective against all classes of viruses, gram positive and gram negative bacteria and fungi. It is odourless, tasteless and bio-degradable. Effective in hard water and organic soiling.

INSTRUCTIONS FOR USE

Following removal of litter and pre-cleaning of house floors, walls and equipment use at a rate of 50gms per 15 litres of water (0.3% solution, 1 in 300 dilution) Use either a pressure washer at a low setting or knapsack sprayer with a wide angle spray. Cover all surfaces with the solution, roof spaces, walls and all equipment.

AERIAL DISINFECTION

Dilute 1 part Hemexol Virucidal powder in 250 parts water, apply as a fine mist over livestock daily using a pressure washer or mechanical fogger.

FOOT AND WHEEL DIPS

Apply 1 part Hemexol Virucidal powder to 150 parts water to all foot and wheel dips. Replace frequently especially if heavy soiling occurs.

FOR USE AGAINST VIRUSES

For effective use against viruses (except Parvovirus) fungi, yeast moulds. Dilute 1 in 100 (1% solution) For control of Parvovirus dilute 1 in 50 (2% solution)

AQUACULTURE

For effective control of viral, bacterial and fungal disease dilute 1 in 100 and soak equipment including boots, waders, dive suit hoses brushes, nets etc., for 30 minutes, in the solution of Hemexol Virucidal powder followed by rinsing in clean water

WATER SANITATION

During terminal disinfection drain header tank and clean, fill tank and dose with Hemexol Virucidal powder at a dilution of 1 in 250. Allow solution to stand in the system for 1 hour, drain and flush with clean water. With birds in place dose the header tank on a regular basis with a 1 in 1000 dilution

DEFRA APPROVAL

For the following notifiable diseases:-

Diseases of Poultry	1 in 300 dilution
Foot and Mouth disease	1 in 1300 dilution

INDEPENDENTLY TESTED

Escherichia Coli	1 in 1000 dilution
Pseudomonas Aërginosa	1 in 1000 dilution
Salmonella Typhi	1 in 1000 dilution

Canine Hepatitis (Adenoviridae)	1 in 100 dilution
Feline Enteritis (Retroviridae)	1 in 200 dilution
Equine Influenza (Orthomyxoviridae)	1 in 300 dilution
HIV 111	1 in 200 dilution
Pox Viridae	1 in 400 dilution
Newcastle Disease	1 in 400 dilution
Gumboro Disease	1 in 250 dilution
Swine Visicular Disease	1 in 240 dilution
P.R.R.S.V	1 in 300 dilution
C.S.F.V.	1 in 300 dilution
I.P.N.	1 in 150 dilution
African Swine Fever	1 in 800 dilution

(Independent report available Chile)

FARM HYGIENE PROCEDURE CONTINUOUS PROTECTION

FOGGING

On a regular basis and frequently at times of pathogenic challenge fog a solution of Hemexol Virucidal powder over livestock N.B. When removing dead birds sprinkle Hemexol Virucidal powder over the contaminated area.


WATER SYSTEMS

Add Hemexol Virucidal powder to the header tank on a regular or daily basis to maintain a dilution of 1 in 1000. Discontinue before live vaccination.

FOOT AND WHEEL DIPS

Maintain Foot and Wheel dips on a regular basis

FULL BIOSECURITY PROGRAMS AVAILABLE ON REQUEST



The information contained in this technical information note is believed to be accurate but each customer is advised to ensure that the product is suitable for their own purpose before using it.

